

Phantom crossing and quintessence limit in extended nonlinear massive gravity

Emmanuel N. Saridakis^{a,b,c}

^a*Physics Division, National Technical University of Athens, 15780 Zografou Campus, Athens, Greece*

^b*CASPER, Physics Department, Baylor University, Waco, TX 76798-7310, USA*

^c*Institut d'Astrophysique de Paris, UMR 7095-CNRS, Université Pierre & Marie Curie, 98bis boulevard Arago, 75014 Paris, France*

E-mail: Emmanuel.Saridakis@baylor.edu

ABSTRACT: We investigate the cosmological evolution in a universe governed by the extended, varying-mass, nonlinear massive gravity, in which the graviton mass is promoted to a scalar-field. We find that the dynamics always leads the varying graviton mass to zero at late times, offering a natural explanation for its hugely-constrained observed value. Despite the limit of the scenario towards standard quintessence, at early and intermediate times it gives rise to an effective dark energy sector of a dynamical nature, which can also lie in the phantom regime, from which it always exits naturally, escaping a Big-Rip. Interestingly enough, although the motivation of massive gravity is to obtain an IR modification, its varying-mass extension in cosmological frameworks leads rather to early and intermediate times modification, and thus to a UV modification instead.

KEYWORDS: Dark energy, phantom crossing, modified gravity, massive gravity, quintessence

Contents

1	Introduction	1
2	Extended nonlinear massive gravity	2
3	Cosmological equations	2
4	Cosmological behavior	4
5	Discussion	7

1 Introduction

The idea of adding mass to the graviton is quite old [1], but the necessary nonlinear terms [2] that can give rise to continuity of the observables [3, 4] lead also to Boulware-Deser (BD) ghosts [5], making the theory unstable. However, recently, a nonlinear extension of massive gravity has been constructed [6, 7] such that the Boulware-Deser ghost is systematically removed (see [8] for a review). The theoretical and phenomenological advantages, amongst which is the universe self-acceleration arising exactly from this IR gravity modification, brought this theory to a significant attention [9–49].

Despite the successes of massive gravity, in the case where the physical and the fiducial metrics have simple homogeneous and isotropic forms the theory proves to be unstable at the perturbation level [40], which led some authors to start constructing less symmetric models [13, 41]. However, in [50] a different approach was followed, namely to extend the theory in a way that the graviton mass is varying, and this was achieved by introducing an extra scalar field which coupling to the graviton potentials produces an effective, varying, graviton mass.

In this work we desire to explore the cosmological implications of this “extended”, varying-mass, massive gravity. As we show, at least in simple cosmological ansatzes, the dynamics always leads the varying graviton mass to zero at late times, and thus the theory has as a limit the standard quintessence paradigm. However, at intermediate times the varying graviton mass leads to very interesting behavior, with a dynamical effective dark energy sector which can easily lie in the phantom regime. Strictly speaking, although the motivation of massive gravity is to obtain an IR modification, its extension in cosmological frameworks leads rather to early and intermediate times modification, and thus to a radical UV modification instead.

2 Extended nonlinear massive gravity

Let us briefly review the “mass-varying massive gravity” that was recently presented in [50]. Their construction is based on the promotion of the graviton mass to a scalar-field function (potential), with the additional insertion in the action of this scalar field’s kinetic term and standard potential. Since such a modification is deeper than allowing for a varying mass, we prefer to call it “extended” nonlinear massive gravity.

In such a construction the action writes as

$$S = \int d^4x \sqrt{-g} \left[\frac{M_P^2}{2} R + V(\psi)(U_2 + \alpha_3 U_3 + \alpha_4 U_4) - \frac{1}{2} \partial_\mu \psi \partial^\mu \psi - W(\psi) \right], \quad (2.1)$$

where M_P is the Planck mass, R the Ricci scalar, and ψ is the new scalar field with $W(\psi)$ its standard potential and $V(\psi)$ its coupling potential which spontaneously breaks general covariance. Furthermore, as usual α_3 and α_4 are dimensionless parameters, and the graviton potentials are given by

$$U_2 = \mathcal{K}_{[\mu}^\mu \mathcal{K}_{\nu]}^\nu, \quad U_3 = \mathcal{K}_{[\mu}^\mu \mathcal{K}_{\nu]}^\nu \mathcal{K}_\rho^\rho, \quad U_4 = \mathcal{K}_{[\mu}^\mu \mathcal{K}_{\nu]}^\nu \mathcal{K}_\rho^\rho \mathcal{K}_\sigma^\sigma, \quad (2.2)$$

with $\mathcal{K}_{[\mu}^\mu \mathcal{K}_{\nu]}^\nu = (\mathcal{K}_\mu^\mu \mathcal{K}_\nu^\nu - \mathcal{K}_\nu^\mu \mathcal{K}_\mu^\nu)/2$ and similarly for the other antisymmetric expressions, and

$$\mathcal{K}_\nu^\mu = \delta_\nu^\mu - \sqrt{g^{\mu\rho} f_{AB} \partial_\rho \phi^A \partial_\nu \phi^B}. \quad (2.3)$$

As in standard massive gravity f_{AB} is a fiducial metric, and the four $\phi^A(x)$ are the Stückelberg scalars introduced to restore general covariance [51], and in the particular case where the f_{AB} is the Minkowski metric they form Lorentz 4-vectors in the internal space and the theory presents a global Poincaré symmetry, too. Finally, one can show that the above extended massive gravity is still free of the Boulware-Deser ghost [50].

3 Cosmological equations

Let us now examine cosmological scenarios in a universe governed by the extended nonlinear massive gravity. For simplicity we consider the fiducial metric to be Minkowski¹

$$f_{AB} = \eta_{AB}, \quad (3.1)$$

and without loss of generality we assume that the dynamical and fiducial metrics are diagonalized simultaneously. For the physical metric we consider for simplicity a flat Friedmann-Robertson-Walker (FRW) form, although one can straightforwardly examine the non-flat case too [50]:

$$d^2 s = -N(\tau)^2 d\tau^2 + a(\tau)^2 \delta_{ij} dx^i dx^j, \quad (3.2)$$

with $N(\tau)$ the lapse function and $a(\tau)$ the scale factor, and for simplicity for the Stückelberg fields we choose the ansatz

$$\phi^0 = b(\tau), \quad \phi^i = x^i. \quad (3.3)$$

¹Note that this case includes the subclasses where f_{AB} can be brought to the Minkowski metric by general coordinate transformation, as we can always choose a gauge for the Stückelberg fields ϕ^A [50].

Finally, in order to obtain a realistic cosmology one includes the usual matter action S_m , coupled minimally to the dynamical metric, corresponding to energy density ρ_m and pressure p_m . Although the above specific application is only a simple subclass of the rich set of possible scenarios, it proves to exhibit very interesting cosmological behavior.

Variation of the total action $S+S_m$ with respect to N and a provides the two Friedmann equations

$$3M_P^2 H^2 = \rho_{DE} + \rho_m, \quad (3.4)$$

$$-2M_P^2 \dot{H} = \rho_{DE} + p_{DE} + \rho_m + p_m, \quad (3.5)$$

where we have defined the Hubble parameter $H = \dot{a}/a$, with $\dot{a} = da/(Nd\tau)$, and in the end we set $N = 1$. In the above expressions we have defined the energy density and pressure of the effective dark energy sector as

$$\rho_{DE} = \frac{1}{2}\dot{\psi}^2 + W(\psi) + V(\psi) \left(\frac{1}{a} - 1 \right) [f_3(a) + f_1(a)] \quad (3.6)$$

$$p_{DE} = \frac{1}{2}\dot{\psi}^2 - W(\psi) - V(\psi)f_4(a) - V(\psi)\dot{b}f_1(a), \quad (3.7)$$

having also introduced the convenient functions

$$\begin{aligned} f_1(a) &= 3 - \frac{2}{a} + \alpha_3 \left(3 - \frac{1}{a} \right) \left(1 - \frac{1}{a} \right) + \alpha_4 \left(1 - \frac{1}{a} \right)^2 \\ f_2(a) &= 1 - \frac{1}{a} + \alpha_3 \left(1 - \frac{1}{a} \right)^2 + \frac{\alpha_4}{3} \left(1 - \frac{1}{a} \right)^3 \\ f_3(a) &= 3 - \frac{1}{a} + \alpha_3 \left(1 - \frac{1}{a} \right) \\ f_4(a) &= - \left[6 - \frac{6}{a} + \left(\frac{1}{a} \right)^2 + \alpha_3 \left(1 - \frac{1}{a} \right) \left(4 - \frac{2}{a} \right) + \alpha_4 \left(1 - \frac{1}{a} \right)^2 \right]. \end{aligned} \quad (3.8)$$

Note that there exists a $\frac{1}{a}$ factor in the expressions of dark energy density and pressure, as well as in the functions f_1 , f_2 , f_3 and f_4 . The value 1 is the default reference that we have chosen in the fixed physical metric of form (3.2) and the Stückelberg fields of form (3.3). One can set the reference value to an arbitrary one a_{ref} , by rescaling a by a/a_{ref} in the metric or transforming ϕ^i to ϕ^i/a_{ref} in the Stückelberg fields, but for simplicity throughout this work we set $a_{ref} = 1$.

One can easily verify that the dark energy density and pressure satisfy the usual evolution equation

$$\dot{\rho}_{DE} + 3H(\rho_{DE} + p_{DE}) = 0, \quad (3.9)$$

and we can also define the dark-energy equation-of-state parameter as usual as

$$w_{DE} \equiv \frac{p_{DE}}{\rho_{DE}}. \quad (3.10)$$

Note that in [50] the authors had named the aforementioned “dark energy” sector as “massive gravity” one, and the quantities ρ_{DE} and p_{DE} as ρ_{MG} and p_{MG} . However, since in this work we focus to late time cosmological behavior, we prefer the above name.

Variation of the total action $S + S_m$ with respect to ψ provides the scalar-field evolution equation:

$$\ddot{\psi} + 3H\dot{\psi} + \frac{dW}{d\psi} + \frac{dV}{d\psi} \left[\left(\frac{1}{a} - 1 \right) [f_3(a) + f_1(a)] + 3\dot{b}f_2(a) \right] = 0. \quad (3.11)$$

Furthermore, variation of $S + S_m$ with respect to b provides the constrain equation

$$V(\psi)Hf_1(a) + \dot{V}(\psi)f_2(a) = 0. \quad (3.12)$$

Finally, one can also extract the matter evolution equation $\dot{\rho}_m + 3H(\rho_m + p_m) = 0$.

4 Cosmological behavior

The cosmological implications of extended nonlinear massive gravity, namely equations (3.4), (3.5), (3.11) and (3.12), prove to be very interesting, however, at least in its present simple but general example, it can be radically different than the usual massive gravity. The reason for this is the constraint equation (3.12), which dramatically restricts $V(\psi)$. Note that in standard massive gravity this equation imposes strong constraints on $b(\tau)$ [13, 16, 49], but in the present model the extra freedom brings these constraints on $V(\psi)$ instead, leaving $b(\tau)$ free. In particular, for $f_2(a) \neq 0$ (which is the case in general) we obtain

$$V(\psi(\tau)) = V_0 e^{-\int \frac{f_1}{af_2} da} = \frac{V_0}{(a-1)[\alpha_4 - (3\alpha_3 + 2\alpha_4)a + (3 + 3\alpha_3 + \alpha_4)a^2]}, \quad (4.1)$$

where we have used the definitions (3.8). The importance of the constraint equation (3.12) was not revealed in [50], where the authors claimed it to be a just a requirement for $V(\psi)$ ². However, from its solution (4.1) we can immediately observe that in general at late times $V(\psi(\tau))$, that is the graviton mass, always goes to zero, independently of the details of the model such are the values of α_3, α_4 , the potential $W(\psi)$, and the matter content. This means that the present scenario of extended nonlinear massive gravity, in cosmological frameworks, cannot provide the usual massive gravity³, and on the contrary it always gives the standard gravity along with the standard quintessence scenario [52, 53]. Similarly, once introduced, the scalar-field cannot be set to zero by hand, since this is not a solution of (3.11) and (3.12) (unless we also set $V(\psi) = 0$ but in this case the model coincides completely with standard quintessence), that is ψ will always have a non-trivial dynamics.

However, although at late times the present scenario coincides with standard quintessence, it can have a very interesting behavior at intermediate times. In particular, the dark energy

²In particular, the specific examples that were considered in [50] were exactly those fine-tuned parameter choices that lead to $f_1(a) = f_2(a) = 0$ and thus to a trivial satisfaction of the constraint (3.12).

³In this sense perhaps the name “extended” nonlinear massive gravity is not appropriate.

sector is not only dynamical, but it can easily lie at the phantom regime [54–57]. This can be seen by observing ρ_{DE} and p_{DE} from (3.6),(3.7), which using the constraint equation (3.12) give

$$\rho_{DE} + p_{DE} = \dot{\psi}^2 - V(\psi) \left(\dot{b} - \frac{1}{a} \right) f_1(a). \quad (4.2)$$

So we can always find regions in the α_3, α_4 parameter space, that can lead to $p_{DE} + \rho_{DE} < 0$ at some stage of the evolution (with a potential $W(\psi)$ that will not lead to large $\dot{\psi}$), even if we require to always have $\rho_{DE} > 0$ (which does not need to be the case in general). This null energy condition violation is always canceled at late times, where the vanishing of the graviton mass leads to $w_{DE} \geq -1$.

In order to present the above behavior in a more transparent way, we numerically evolve the scenario at hand choosing without loss of generality $b(\tau) = \ln[a(\tau)]$ (with the dimensional coefficient set to 1 in M_P^2 -units), for dust matter ($w_m \equiv p_m/\rho_m = 0$), using the redshift $z = a_0/a - 1$ as the independent variable (with a_0 the present scale factor), and imposing the present dark energy density $\Omega_{DE} \equiv \rho_{DE}/(3M_P^2 H^2)$ to be ≈ 0.72 and its initial value to be ≈ 0 . In Fig. 1 we present the effective dark-energy equation-of-state parameter w_{DE} as a function of z , in the case where the usual scalar field potential is $W(\psi) = W_0 = \text{const.}$, and for two choices of the parameters $\alpha_3, \alpha_4, W_0, V_0$ ⁴. As described




Figure 1. Evolution of the dark energy equation-of-state parameter w_{DE} as a function of the redshift z , in the case where the usual scalar field potential is $W(\psi) = W_0 = \text{const.}$ The black-solid curve corresponds to $\alpha_3 = -1, \alpha_4 = -1, W_0 = 1, V_0 = 40$, while the red-dashed curve corresponds to $\alpha_3 = -2, \alpha_4 = -2, W_0 = 2.5, V_0 = 6$, in M_P^2 -units. The dotted -1-line is depicted for convenience.

⁴Note that the graviton mass is proportional to V_0 but significantly downgraded by the denominator of (4.1), thus it is below the Planck mass even if V_0 is chosen larger than M_P^4 .

above, at early and intermediate times the coupling potential $V(\psi)$ is non-zero leading w_{DE} to exhibit a dynamical nature, which can lie in the quintessence regime (black-solid curve) or in the phantom regime (red-dashed curve). Additionally, as expected, since in both cases the usual scalar potential is constant, the system at late times, where the coupling potential $V(\psi)$ of (4.1) becomes zero, tends to the cosmological constant behavior.

In fig. 2 we present w_{DE} as a function of z , in the case where the usual scalar field potential has the exponential form $W(\psi) = W_0 e^{-\lambda\psi}$, and for three choices of the parameters $\alpha_3, \alpha_4, \lambda, W_0, V_0$. In all three sub-cases, w_{DE} presents a dynamical nature, which can be non-




Figure 2. Evolution of the dark energy equation-of-state parameter w_{DE} as a function of the redshift z , in the case where the usual scalar field potential takes the exponential form $W(\psi) = W_0 e^{-\lambda\psi}$. The black-solid curve corresponds to $\alpha_3 = 2$, $\alpha_4 = 2$, $\lambda = 0.5$, $W_0 = 15$, $V_0 = 12$, the red-dashed curve corresponds to $\alpha_3 = 1$, $\alpha_4 = 1$, $\lambda = 0.5$, $W_0 = 5$, $V_0 = 2.5$, and the green-dotted curve corresponds to $\alpha_3 = -2$, $\alpha_4 = -2$, $\lambda = 1$, $W_0 = 1$, $V_0 = 2$, in M_P^2 -units. The dotted -1-line is depicted for convenience. In order to depict the limiting behavior towards quintessence, we have evolved the system towards the future and up to $z = -0.9$.

monotonic or/and it can cross the phantom divide from below to above [58]. However, as described above, at late times, where the coupling $V(\psi)$ becomes zero, all three sub-cases are led to their usual quintessence limit, where the final w_{DE} is determined solely from the W -potential exponent λ [59], and such a limiting behavior can be clearly seen if we evolve the system towards the future and up to $z = -0.9$.

In summary, as we can see the scenario at hand exhibits very interesting cosmological behavior at early and intermediate times, with a dynamical dark energy sector which can additionally lie in the phantom regime, before limit towards the standard quintessence scenario. Note that despite the phantom-crossing, at late times we always obtain $w_{DE} \geq -1$ since the vanishing of the graviton mass restores the null energy condition for the effective

dark energy sector, that is the universe will always escape from the phantom regime and the Big-Rip future [60, 61] that is common to the majority of phantom models.

5 Discussion

In this work we investigated the cosmological evolution in a universe governed by the extended, varying-mass, nonlinear massive gravity. Even for simple ansatzes the scenario proves to have a very interesting behavior, comparing with standard massive gravity.

The first result is that the dynamics in cosmological frameworks always leads the varying graviton mass to zero at late times, and thus the theory possesses as a limit the standard quintessence paradigm. This is a great advantage of the present construction, since it offers a natural explanation of the tiny and hugely-constrained graviton mass that arises from current observations. The graviton mass does not have to be tuned to an amazingly small number, as it is the case in standard massive gravity, but it is the dynamics that lead it asymptotically to zero.

Despite the vanishing of the graviton mass at late times, and the limit of the scenario towards standard quintessence, at early and intermediate ones it can lead to very interesting behavior. In particular, it can give rise to an effective dark energy sector of a dynamical nature, which can also lie in the phantom regime. The violation of the null energy condition for the effective dark energy sector at intermediate times arises naturally for suitable (not fine-tuned) regions in the Lagrangian parameters, and it is always canceled at late times due to the vanishing of the graviton mass. These features are in agreement with observations and they offer an explanation for the dynamical evolution of the dark-energy equation-of-state parameter, for its relaxation close or at the cosmological constant value, and also for the indicated possibility to have crossed the phantom divide. Moreover, even if it enters the phantom regime, the scenario at hand always returns naturally to the quintessence one, offering a solution to the Big-Rip fate of the standard phantom scenarios.

We mention here that although we performed the above analysis in a flat universe, and with the fiducial metric to be Minkowski, qualitatively the obtained behavior is not a result of them (one can easily see that they hold in the non-flat case of [50] too) but it arises from the deeper structure of the theory, namely from the scalar-field coupling to the graviton potential. Thus, we do not expect the results to change in more general cases, unless one fine-tunes the theory.

In conclusion, the extended, varying-mass, nonlinear massive gravity leads to very interesting cosmological behavior at early and intermediate times, while it always limits towards the standard quintessence scenario, where the graviton is massless and the extra scalar is only minimally coupled to gravity. Strictly speaking, although the motivation of massive gravity is to obtain an IR modification, its varying-mass extension in cosmological frameworks leads rather to early and intermediate times modification, and thus to a UV modification instead.

Acknowledgments

We wish to thank Y-F. Cai, A. Gumrukcuoglu, Q-G. Huang, T. Koivisto, C. Lin, S. Mukohyama, Y.-S. Piao, N. Tamanini and S.-Y. Zhou, for useful comments. The research project is implemented within the framework of the Action Supporting Postdoctoral Researchers of the Operational Program “Education and Lifelong Learning” (Actions Beneficiary: General Secretariat for Research and Technology), and is co-financed by the European Social Fund (ESF) and the Greek State.

References

- [1] M. Fierz, W. Pauli, *On relativistic wave equations for particles of arbitrary spin in an electromagnetic field*, Proc. Roy. Soc. Lond. **A173**, 211 (1939).
- [2] A. I. Vainshtein, *To the problem of nonvanishing gravitation mass*, Phys. Lett. B **39**, 393 (1972).
- [3] H. van Dam, M. J. G. Veltman, *Massive and massless Yang-Mills and gravitational fields*, Nucl. Phys. **B22**, 397 (1970).
- [4] V. I. Zakharov, *Linearized gravitation theory and the graviton mass*, JETP Lett. **12**, 312 (1970).
- [5] D. G. Boulware, S. Deser, *Can gravitation have a finite range?*, Phys. Rev. **D6**, 3368 (1972).
- [6] C. de Rham, G. Gabadadze, *Generalization of the Fierz-Pauli Action*, Phys. Rev. **D82**, 044020 (2010), [[arXiv:1007.0443](#)].
- [7] C. de Rham, G. Gabadadze and A. J. Tolley, *Resummation of Massive Gravity*, Phys. Rev. Lett. **106**, 231101 (2011), [[arXiv:1011.1232](#)].
- [8] K. Hinterbichler, *Theoretical Aspects of Massive Gravity*, Rev. Mod. Phys. **84**, 671 (2012), [[arXiv:1105.3735](#)].
- [9] K. Koyama, G. Niz and G. Tasinato, *Strong interactions and exact solutions in non-linear massive gravity*, Phys. Rev. D **84** (2011) 064033, [[arXiv:1104.2143](#)].
- [10] S. F. Hassan and R. A. Rosen, *Resolving the Ghost Problem in non-Linear Massive Gravity*, Phys. Rev. Lett. **108**, 041101 (2012), [[arXiv:1106.3344](#)].
- [11] C. de Rham, G. Gabadadze and A. Tolley, *Ghost free Massive Gravity in the Stückelberg language*, Phys. Lett. B **711**, 190 (2012), [[arXiv:1107.3820](#)].
- [12] B. Cuadros-Melgar, E. Papantonopoulos, M. Tsoukalas and V. Zamarias, *Massive Gravity with Anisotropic Scaling*, [[arXiv:1108.3771](#)].
- [13] G. D’Amico, C. de Rham, S. Dubovsky, G. Gabadadze, D. Pirtskhalava, A. J. Tolley, *Massive Cosmologies*, [[arXiv:1108.5231](#)].
- [14] S. F. Hassan and R. A. Rosen, *Bimetric Gravity from Ghost-free Massive Gravity*, JHEP **1202**, 126 (2012), [[arXiv:1109.3515](#)].
- [15] J. Kluson, *Note About Hamiltonian Structure of Non-Linear Massive Gravity*, JHEP **1201**, 013 (2012), [[arXiv:1109.3052](#)].
- [16] A. E. Gumrukcuoglu, C. Lin and S. Mukohyama, *Open FRW universes and self-acceleration from nonlinear massive gravity*, JCAP **1111**, 030 (2011), [[arXiv:1109.3845](#)].

- [17] M. S. Volkov, *Cosmological solutions with massive gravitons in the bigravity theory*, JHEP **1201**, 035 (2012), [[arXiv:1110.6153](#)].
- [18] M. von Strauss, A. Schmidt-May, J. Enander, E. Mortsell and S. F. Hassan, *Cosmological Solutions in Bimetric Gravity and their Observational Tests*, JCAP **1203**, 042 (2012), [[arXiv:1111.1655](#)].
- [19] D. Comelli, M. Crisostomi, F. Nesti and L. Pilo, *FRW Cosmology in Ghost Free Massive Gravity*, JHEP **1203**, 067 (2012) [Erratum-ibid. **1206**, 020 (2012)], [[arXiv:1111.1983](#)].
- [20] S. F. Hassan and R. A. Rosen, *Confirmation of the Secondary Constraint and Absence of Ghost in Massive Gravity and Bimetric Gravity*, JHEP **1204**, 123 (2012), [[arXiv:1111.2070](#)].
- [21] L. Berezhiani, G. Chkareuli, C. de Rham, G. Gabadadze and A. J. Tolley, *On Black Holes in Massive Gravity*, Phys. Rev. D **85**, 044024 (2012), [[arXiv:1111.3613](#)].
- [22] A. E. Gumrukcuoglu, C. Lin and S. Mukohyama, *Cosmological perturbations of self-accelerating universe in nonlinear massive gravity*, JCAP **1203**, 006 (2012), [[arXiv:1111.4107](#)].
- [23] N. Khosravi, N. Rahmanpour, H. R. Sepangi and S. Shahidi, *Multi-Metric Gravity via Massive Gravity*, Phys. Rev. D **85**, 024049 (2012), [[arXiv:1111.5346](#)].
- [24] Y. Brihaye and Y. Verbin, *Perfect Fluid Spherically-Symmetric Solutions in Massive Gravity*, [[arXiv:1112.1901](#)].
- [25] I. L. Buchbinder, D. D. Pereira and I. L. Shapiro, *One-loop divergences in massive gravity theory*, Phys. Lett. B **712**, 104 (2012), [[arXiv:1201.3145](#)].
- [26] H. Ahmedov and A. N. Aliev, *Type N Spacetimes as Solutions of Extended New Massive Gravity*, Phys. Lett. B **711**, 117 (2012), [[arXiv:1201.5724](#)].
- [27] E. A. Bergshoeff, J. J. Fernandez-Melgarejo, J. Rosseel and P. K. Townsend, *On 'New Massive' 4D Gravity*, JHEP **1204**, 070 (2012), [[arXiv:1202.1501](#)].
- [28] M. Crisostomi, D. Comelli and L. Pilo, *Perturbations in Massive Gravity Cosmology*, [[arXiv:1202.1986](#)].
- [29] M. F. Paulos and A. J. Tolley, *Massive Gravity Theories and limits of Ghost-free Bigravity models*, [[arXiv:1203.4268](#)].
- [30] S. F. Hassan, A. Schmidt-May and M. von Strauss, *Proof of Consistency of Nonlinear Massive Gravity in the Stückelberg Formulation*, [[arXiv:1203.5283](#)].
- [31] D. Comelli, M. Crisostomi, F. Nesti and L. Pilo, *Degrees of Freedom in Massive Gravity*, [[arXiv:1204.1027](#)].
- [32] F. Sbisà, G. Niz, K. Koyama and G. Tasinato, *Characterising Vainshtein Solutions in Massive Gravity*, [[arXiv:1204.1193](#)].
- [33] J. Kluson, *Non-Linear Massive Gravity with Additional Primary Constraint and Absence of Ghosts*, [[arXiv:1204.2957](#)].
- [34] G. Tasinato, K. Koyama and G. Niz, *New symmetries in Fierz-Pauli massive gravity*, [[arXiv:1204.5880](#)].
- [35] K. Morand and S. N. Solodukhin, *Dual Massive Gravity*, [[arXiv:1204.6224](#)].
- [36] V. F. Cardone, N. Radicella and L. Parisi, *Constraining massive gravity with recent*

- cosmological data*, Phys. Rev. D **85**, 124005 (2012), [[arXiv:1205.1613](#)].
- [37] V. Baccetti, P. Martin-Moruno and M. Visser, *Massive gravity from bimetric gravity*, [[arXiv:1205.2158](#)].
 - [38] P. Gratia, W. Hu and M. Wyman, *Self-accelerating Massive Gravity: Exact solutions for any isotropic matter distribution*, [[arXiv:1205.4241](#)].
 - [39] M. S. Volkov, *Exact self-accelerating cosmologies in the ghost-free bigravity and massive gravity*, [[arXiv:1205.5713](#)].
 - [40] A. De Felice, A. E. Gumrukcuoglu and S. Mukohyama, *Massive gravity: nonlinear instability of the homogeneous and isotropic universe*, [[arXiv:1206.2080](#)].
 - [41] A. E. Gumrukcuoglu, C. Lin and S. Mukohyama, *Anisotropic Friedmann-Robertson-Walker universe from nonlinear massive gravity*, [[arXiv:1206.2723](#)].
 - [42] C. de Rham and S. Renaux-Petel, *Massive Gravity on de Sitter and Unique Candidate for Partially Massless Gravity*, [[arXiv:1206.3482](#)].
 - [43] M. Berg, I. Buchberger, J. Enander, E. Mortsell and S. Sjors, *Growth Histories in Bimetric Massive Gravity*, [[arXiv:1206.3496](#)].
 - [44] G. D’Amico, *Cosmology and perturbations in massive gravity*, [[arXiv:1206.3617](#)].
 - [45] V. Baccetti, P. Martin-Moruno and M. Visser, *Null Energy Condition violations in bimetric gravity*, [[arXiv:1206.3814](#)].
 - [46] M. Fasiello and A. J. Tolley, *Cosmological perturbations in Massive Gravity and the Higuchi bound*, [[arXiv:1206.3852](#)].
 - [47] G. D’Amico, G. Gabadadze, L. Hui and D. Pirtskhalava, *Quasi-Dilaton: Theory and Cosmology*, [[arXiv:1206.4253](#)].
 - [48] V. Baccetti, P. Martin-Moruno and M. Visser, *Gordon and Kerr-Schild ansatze in massive and bimetric gravity*, [[arXiv:1206.4720](#)].
 - [49] D. Langlois and A. Naruko, *Cosmological solutions of massive gravity on de Sitter*, [[arXiv:1206.6810](#)].
 - [50] Q. -G. Huang, Y. -S. Piao and S. -Y. Zhou, *Mass-Varying Massive Gravity*, [[arXiv:1206.5678](#)].
 - [51] N. Arkani-Hamed, H. Georgi and M. D. Schwartz, *Effective field theory for massive gravitons and gravity in theory space*, Annals Phys. **305**, 96 (2003), [[arXiv:hep-th/0210184](#)].
 - [52] P. J. E. Peebles and B. Ratra, *Cosmology with a Time Variable Cosmological Constant*, Astrophys. J. **325**, L17 (1988).
 - [53] C. Wetterich, *Cosmology and the Fate of Dilatation Symmetry*, Nucl. Phys. B **302**, 668 (1988).
 - [54] R. R. Caldwell, *A Phantom menace?*, Phys. Lett. B **545**, 23 (2002), [[arXiv:astro-ph/9908168](#)].
 - [55] X. -m. Chen, Y. -g. Gong and E. N. Saridakis, *Phase-space analysis of interacting phantom cosmology*, JCAP **0904**, 001 (2009), [[arXiv:0812.1117](#)].
 - [56] E. N. Saridakis, *Phantom evolution in power-law potentials*, Nucl. Phys. B **819**, 116 (2009), [[arXiv:0902.3978](#)].

- [57] E. N. Saridakis and S. V. Sushkov, *Quintessence and phantom cosmology with non-minimal derivative coupling*, Phys. Rev. D **81**, 083510 (2010), [[arXiv:1002.3478](#)].
- [58] Y. -F. Cai, E. N. Saridakis, M. R. Setare and J. -Q. Xia, *Quintom Cosmology: Theoretical implications and observations*, Phys. Rept. **493**, 1 (2010), [[arXiv:0909.2776](#)].
- [59] E. J. Copeland, A. R. Liddle and D. Wands, *Exponential potentials and cosmological scaling solutions*, Phys. Rev. D **57**, 4686 (1998), [[arXiv:gr-qc/9711068](#)].
- [60] F. Briscese, E. Elizalde, S. Nojiri and S. D. Odintsov, *Phantom scalar dark energy as modified gravity: Understanding the origin of the Big Rip singularity*, Phys. Lett. B **646**, 105 (2007), [[arXiv:hep-th/0612220](#)].
- [61] S. Capozziello, M. De Laurentis, S. Nojiri and S. D. Odintsov, *Classifying and avoiding singularities in the alternative gravity dark energy models*, Phys. Rev. D **79**, 124007 (2009), [[arXiv:0903.2753](#)].